Contents to volume 241

Hypothesis

Do mitochondrial DNA fragments promote cancer and aging? (FEB 06508) *C. Richter* 1

Research Letters

Interleukin 1 and tumour necrosis factor increase phosphorylation of fibroblast proteins (FEB 06539) *P. Kaur, J. Saklatvala* 6

Heparan sulphate with no affinity for antithrombin III and the control of haemostasis (FEB 06538) *M.F. Scully, V. Ellis, V.V. Kakkar* 11

The influence of culture medium composition on drug metabolising enzyme activities of the human liver derived Hep G2 cell line (FEB 06540) *H. Doostdar, S.J. Duthie, M.D. Burke, W.T. Melvin, M.H. Grant* 15

Programmed cell death: dying cells synthesize a co-ordinated, unique set of proteins in two different episodes of cell death (FEB 06505) *A.G. Wadewitz, R.A. Lockshin* 19

A biphasic nature of the bleomycin-mediated degradation of DNA (FEB 06537) *L. Strekowski, J.L. Mokrosz, W.D. Wilson* 24

Rapid preparation of the nicotinic acetylcholine receptor for crystallization in detergent solution (FEB 06542) *S. Hertling-Jaweed, G. Bandini, A. Müller-Fahrnow, V. Dommes, F. Hucho* 29

Spectroscopic studies on the mode of binding of ATP, UTP and α-amanitin with yeast RNA polymerase II (FEB 06536) *P. Bhargava, D. Chatterji* 33

Conformation of 1,4-dihydropyridine – planar or boat-like? (FEB 06527) *H.-J. Hofmann, R. Cimiraglia* 38

The complete amino acid sequence of bovine milk angiogenin (FEB 06534) *P. Maes, D. Damart, C. Rommens, J. Montreuil, G. Spik, A. Tartar* 41

Engineering of an intersubunit disulphide bridge in glutathione reductase from *Escherichia coli* (FEB 06526) *N.S. Scrutton, A. Berry, R.N. Perham* 46

Allosteric inhibition of *Dictyostelium discoideum* fructose-1,6-bisphosphatase by fructose 2,6-bisphosphate (FEB 06535) *V. Andrés, L. García-Salguero, M.-E. Gómez, J.J. Aragón* 51

Irreversible inactivation of *Saccharomyces cerevisiae* fructose-1,6-bisphosphatase independent of protein phosphorylation at Ser^{11} (FEB 06547) *M. Rose, K.-D. Entian, L. Hofmann, R.F. Vogel, D. Mecke* 55

A methyl-CoM methylreductase system from methanogenic bacterium strain Göl not requiring ATP for activity (FEB 06543) *U. Deppenmeier, M. Blaut, A. Jussofie, G. Gottschalk* 60

A Ca^{2+}-stimulated, Mg^{2+}-dependent ATPase activity in subcellular fractions from *Schistosoma mansoni* (FEB 06554) *V.M.N. Cunha, W. de Souza, F. Noël* 65

Activation of human pepsinogens (FEB 06551) *B. Foltmann* 69

Characterization of mouse reticulocyte free globin mRNP (FEB 06550) *A. Rairkar, R.E. Lockard* 73

Secondary structure of the promastigote surface protease of *Leishmania* (FEB 06541) *F. Jähnig, R. Etges* 79

The *Saccharomyces cerevisiae ARO1* gene: An example of the co-ordinate regulation of five enzymes on a single biosynthetic pathway (FEB 06552) *K. Duncan, R.M. Edwards, J.R. Coggins* 83

Heterogeneous expression of glutamine synthetase mRNA in rat liver parenchyma revealed by in situ hybridization and Northern blot analysis of RNA from periportal and perivenous hepatocytes (FEB 06545) *R. Gebhardt, A. Ebert, G. Bauer* 89

The amyloid precursor protein of Alzheimer disease is expressed as a 130 kDa polypeptide in various cultured cell types (FEB 06546) *L. Autilio-Gambetti, A. Morandi, M. Tabaton, B. Schaetzle, D. Kovacs, G. Perry, B. Greenberg, P. Gambetti* 94

A new photosystem II reaction center component (4.8 kDa protein) encoded by chloroplast genome (FEB 06544) *M. Ikeuchi, Y. Inoue* 99

Published by Elsevier Science Publishers B.V. (Biomedical Division)
00145793/88/$3.50 © 1988 Federation of European Biochemical Societies

The mitochondrial outer membrane channel, VDAC, is modulated by a soluble protein (FEB 06549) *M.J. Holden, M. Colombini* 105
Genetic fusion of a non-toxic heat-stable enterotoxin-related decapeptide antigen to cholera toxin B-subunit (FEB 06553) *J. Sanchez, A.-M. Svennerholm, J. Holmgren* 110
Increased protein synthesis response to insulin in fibroblasts treated with the diacylglycerol kinase inhibitor R59022 (FEB 06558) *J.E. Hesketh, N. McKenzie, G.P. Campbell* 115
Location of a region of the muscarinic acetylcholine receptor involved in selective effector coupling (FEB 06563) *T. Kubo, H. Bujo, I. Akiba, J. Nakai, M. Mishina, S. Numa* 119
A case of carnitine palmitoyltransferase II deficiency in human skeletal muscle (FEB 06557) *R. Singh, I.M. Shepherd, J.P. Derrick, R.R. Ramsay, H.S.A. Sherratt, D.M. Turnbull* 126
Identification of two distinct protein carboxyl methyltransferases in eucaryotic cells (FEB 06559) *R. Solomon, O. Katzir, Y. Egozi, Y. Kloog* 131
Boar proacrosin is a single-chain molecule which has the N-terminus of the acrosin A-chain (light chain) (FEB 06562) *D. Čechová, E. Töpfer-Petersen, A. Henschen* 136
Flagellin parts acquiring a regular structure during polymerization are disposed on the molecule ends (FEB 06567) *A.S. Kostyukova, M.G. Pyatibratov, V.V. Filimonov, O.V. Fedorov* 141
Architectonics of a bacterial flagellin filament subunit (FEB 06568) *O.V. Fedorov, A.S. Kostyukova, M.G. Pyatibratov* 145
Phorbol 12-myristate 13-acetate modulates the cAMP-induced light-scattering response of a *Dictyostelium discoïdeum* cell population (FEB 06555) *R. Thiery, R. Klein, I. Tatischeff* 149
Retinoic acid inhibits phospholipid turnover and protein kinase C activity in RA-sensitive but not in RA-resistant cells (FEB 06560) *M.C. Vanier, D. Banerjee, B.B. Mukherjee* 154
Metallothionein RNA levels in HL-60 cells: Effect of cadmium, differentiation, and protein kinase C activation (FEB 06561) *T. Hanke, M. Tyers, C.B. Harley* 159
Serum induces the immediate opening of Ca^{2+}-activated channels in quiescent human fibroblasts (FEB 06565) *A. Peres, R. Zippel, E. Sturani* 164
Ionic currents in avian granulosa cells (FEB 06548) *J.-L. Schwartz, G.A.R. Mealing, E.K. Asem, J.F. Whitfield, B.K. Tsang* 169
Transcellular ionic currents studied by intracellular potential recordings in *Neurospora crassa* hyphae: Transfer of energy from proximal to apical cells (FEB 06566) *T.V. Potapova, K.B. Aslanidi, T.A. Belozerskaya, N.N. Levina* 173
Calcium mobilization by angiotensin II and neurotransmitter receptors expressed in *Xenopus laevis* oocytes (FEB 06586) *K. Sandberg, A.J. Markwick, D.P. Trinh, K.J. Catt* 177
Apomyoglobin forms a micellar complex with phospholipid at low pH (FEB 06574) *J.W. Lee, H. Kim* 181
Adhesion defect of ascites cells corrected with membrane-bound attachment molecules (FEB 06572) *L.V. Gordeeva, A.A. Bogdanov, jr, B.A. Baibakov, V.P. Torchilin, L.B. Margolis* 185
Effects of ATP depletion on the mechanism of hexose transport in intact human erythrocytes (FEB 06575) *J.M. May* 188
The synthesis of human α-2-HS glycoprotein is down-regulated by cytokines in hepatoma HepG2 cells (FEB 06581) *M. Daveau, C. Davrinche, N. Julen, M. Hiron, P. Arnaud, J.-P. Lebreton* 191
Insulin and progesterone activate a common synthetic ribosomal protein S6 peptide kinase in *Xenopus* oocytes (FEB 06582) *M.F. Cicirelli, S.L. Pelech, E.G. Krebs* 195
Bleomycin-detectable iron in serum from leukaemic patients before and after chemotherapy: Therapeutic implications for treatment with oxidant-generating drugs (FEB 06573) *B. Halliwell, O.I. Aruoma, G. Mufti, A. Bomford* 202
Oxidation of rat insulin II, but not I, leads to anomalous elution profiles upon HPLC analysis of insulin-related peptides (FEB 06583) *D. Gross, A. Skvorak, G. Hendrick, G. Weir, L. Villa-Komaroff, P. Halban* 205
Identification and characterization of glucagon-like peptide-1 7–36 amide-binding sites in the rat brain and lung (FEB 06584) *S.M. Kanse, B. Kreymann, M.A. Ghatei, S.R. Bloom* 209

Partial sequence of the rat heavy neurofilament polypeptide (NF-H): Identification of putative phosphorylation sites (FEB 06569) *K.C. Breen, P.A. Robinson, D. Wion, B.H. Anderton* 213
Cyclic nucleotide phosphodiesterase activity in *Neurospora crassa*: Purification by immunoaffinity chromatography and characterization (FEB 06585) *R.M. Ulloa, C.P. Rubinstein, L. Molina y Vedia, H.N. Torres, M.T. Téllez-Iñón* 219
Protein secondary structure and homology by neural networks: The α-helices in rhodopsin (FEB 06576) *H. Bohr, J. Bohr, S. Brunak, R.M.J. Cotterill, B. Lautrup, L. Nørskov, O.H. Olsen, S.B. Petersen* 223
Identification of *Eco*RV fragments spanning the *N*α-tubulin gene of *Physarum* (FEB 06577) *D.I. Watts, M.J. Monteiro, R.A. Cox* 229
Tetrameric structure of the nonactivated glucocorticoid receptor in cell extracts and intact cells (FEB 06580) *M. Rexin, W. Busch, B. Segnitz, U. Gehring* 234
Electron microscopy and image analysis of the multicatalytic proteinase (FEB 06564) *W. Baumeister, B. Dahlmann, R. Hegerl, F. Kopp, L. Kuehn, G. Pfeifer* 239
Sulfated *N*-linked carbohydrate chains in porcine thyroglobulin (FEB 06523) *J.P. Kamerling, I. Rijkse, A.A.M. Maas, J.A. van Kuik, J.F.G. Vliegenthart* 246
Renaturation of guanidine-unfolded tryptophan synthase by multi-mixing stopped-flow dilution in D_2O (FEB 06571) *S. Blond-Elguindi, B. Friguet, M.E. Goldberg* 251
Evidence of a yeast proteinase specific for elongation factor 2 (FEB 06570) *L. Servillo, L. Quagliuolo, C. Balestrieri, A. Giovane* 257
A novel crystallin from octopus lens (FEB 06556) *S.-H. Chiou* 261

Author Index 265

Tables of Contents 269

Notes to Authors 273

FEBS NEWS

Change of address (Eur. J. Biochem.)/Congress Announcement 275